

Jacobs Journal of Veterinary Science and Research

Research Article

High lungworm burden in enclosed wild boar from Eastern Austria

Bettina Lechner^{*}1, Barbara Hinney, Georg Duscher, Anja Joachim

Institute of Parasitology, Department of Pathobiology Vetmeduni Vienna

^{*}Corresponding author: Mag.med.vet. Bettina Lechner, Institute of Parasitology, Department of Pathobiology, Vetmeduni Vienna

Veterinärplatz 1 A-1210, Vienna, AUSTRIA. Tel: +43 1 25077 2201; Fax: +43 1 25077 2290;

Email: bettina.lechner91@gmail.com

Received: 12-18-2015

Accepted: 02-24-2016

Published: 03-04-2016

Copyright: © 2016 Bettina Lechner

Abstract

Objective

We aimed to investigate the endoparasite status, especially the lungworm burden, of wild boar in an enclosed hunting ground in Eastern Austria.

Animals and samples

In total, 49 lungs and 47 corresponding faecal samples of animals of different age groups (shoats, young pigs and adults) were examined.

Methods

Lungs were dissected and adult metastrongylids were counted and determined to species level (n=25/animal). Faecal samples were prepared by sedimentation/flotation and examined qualitatively and also quantitatively (by McMaster counting).

Results

Adult lungworms were present in 94% of the animals. Eggs of *Metastrongylus* spp. (89%), *Ascaris suum* (8%), *Globocephalus urosubulatus* (96%), *Trichuris suis* (51%), as well as oocysts of coccidia (100%) were detected. Five different species of *Metastrongylus* (*M. apri*, *M. salmi*, *M. pudendotectus*, *M. confusus* and *M. asymmetricus*) were found in the lungs. Oocysts of *E. deblickei*, *E. scabra*, *E. perminuta*, *E. porci*, *E. polita* and *Cystoisospora suis* could be differentiated. *E. deblickei* was the most common (100%), *E. porci* (28%) the rarest species. In most cases, parasite excretion tended to decline with increasing age. In contrast to domestic pigs, *C. suis* could be found in all age groups.

Conclusions

The high intensities and infection rates indicate a considerable infection pressure which may lead to health problems in enclosed wild boar. Proper habitat conservation and hunting practices are necessary to keep a stable and healthy population in such enclosed environments.

Keywords: game animals; nematodes; *Metastrongylus*; *Eimeria*; *Cystoisospora*; *Globocephalus*; *Trichuris*; *Ascaris*

Introduction

Besides roe and red deer, wild boar is one of the most important hoofed game species in Central Europe. Due to the difficult hunt and the excessive food supply the populations are constantly increasing. Growing populations in restricted habitats however pose an increasing risk of parasitic infec-

tions. Especially in game enclosures it must be assumed that the transmission risk for various pathogens increases inordinately [1, 2].

Coccidial infections are frequently asymptomatic, but *Cystoisospora suis* and rarely *Eimeria* spp. can also cause catarrhal, rarely necrotic enteritis with non-haemorrhagic

diarrhoea in young piglets, leading to weight loss and poor performance [3-6].

Little is known about the prevalences in wild boar and the effects on their health. For Austria, so far five species of *Eimeria* have been described from domestic and wild suids. The latter harboured mostly *Eimeria deblickei* and *Eimeria polita*, as well as *C. suis* [7].

The large roundworm *Ascaris suum* can be found in the small intestines of domestic and wild suids and can cause reduced growth when present in large numbers [8]. Similarly, the whipworm *Trichuris suis* frequently resides in the large intestines of wild and domestic pigs and can cause enteritis with diarrhoea in intensive infections, as well as hypalbuminaemia and reduced growth [9]. The porcine hookworm *Globocephalus urosubulatus* on the other hand infects primarily wild boar [10].

The porcine lungworms of the genus *Metastrongylus* are rarely found in domestic pigs since they require earthworms as intermediate hosts so only pigs with outdoor access to earthworm habitats can become infected while wild boars frequently harbour these parasites. Especially in young animals the parasite is rather pathogenic, inducing lung damage with dyspnoea, weight loss and even death [11, 12]. Due their pathogenicity they are still considered to be the most important airway pathogens in extensive pig management [13, 14]. In Europe, so far five different species have been described, *Metastrongylus apri* (syn. *M. elongatus*), *M. salmi*, *M. pudendotectus*, *M. confusus* and *M. asymmetricus* [12, 15-19].

To evaluate the parasite burden of wild boar from an enclosure in Eastern Austria, we investigated lungs and faecal samples of animals from different age categories for infections with parasites, especially lungworms.

Material and Methods


Samples

Samples were provided by hunters and originated from a large enclosure in Eastern Austria. As soon as possible after killing, lungs and faecal samples were kept refrigerated in a cooling chamber at a temperature of 4-8 °C until examination 3-5 days later. The hunters also evaluated age, weight and body condition of the animals. In total 49 animals, 10 male and 23 female shoats (<1 year), two male and six female young animals (1-2 years), four sows and four tuskers were examined. Lungs were available from all of them, faecal samples from all but two shoats (n=47). Lungs were inspected for gross pathological changes, the upper and lower airways were cut open, the lungworms removed and fixed in 70% ethanol.

Lungworm differentiation

Specimens from each positive lung were counted. A maximum of 25 females per each lung were differentiated according to different keys [12, 18, 20]. The species were differentiated according to their characteristic endings shown in Figure 1.

Figure 1. *Metastrongylus* species in wild boar: Posterior ends of females of *M. apri* (A); *M. pudendotectus* (B), *M. asymmetricus* (C), and *M. confusus* (D).



Coproscopy

Using a combined sedimentation/flotation technique, faecal samples were examined for parasite stages (helminth eggs and coccidian oocysts) using saturated sugar solution. Positive samples were examined by a modified McMaster technique [21]. Coccidia oocysts were microscopically differentiated according to their size and morphology [22].

Statistical evaluation

To compare the parasite distribution in different age groups data were analysed by nonparametric tests (Kruskal-Wallis and Mann-Whitney U test) in SPSS v. 20 (SPSS Inc., Chicago, USA). Correlations between body mass and parasite burden/excretion and between different species of *Metastrongylus* were tested calculating Pearson's rank correlation coefficient.

Results and Discussion

Body condition of the animals

The body condition was documented for 48 animals, and 38

of these were in very good or good physical condition. Nine shoats were poorly nourished and a male adult animal was emaciated. For nine of the ten animals with a poor condition faecal samples could be examined. No correlation with the burden of adult lungworms or with the excretion of oocysts/nematode eggs could be established (details not shown).

Lung pathology

During gross examination purulent areas, enlarged lymph nodes, pulmonary oedema and haemorrhage, emphysema and adult lungworms could be seen in different parts of the lung. Haemorrhage was the most common finding, which could at least partly be attributed to gunshot injuries and was not evaluated further. Emphysema was the second most common alteration, found in 90% of the shoats in different parts of the lung. Young pigs showed mainly signs of purulent pneumonia and enlarged lymph nodes (Fig. 1). Earlier works stated that pathological changes induced by porcine lungworms are irreversible [11], indicating the health importance of metastrongylid infections in wild boar.

Metastrongylid infections

In total, 46 animals (94%) harboured adult lungworms. Shoats showed a mean worm burden of 195.5 specimen (SD=251.5,

median 109, maximum number of worms: 1020) which was higher than in the other two groups (Table 1).

Overall, five different species of *Metastrongylus* could be differentiated, *M. apri*, *M. salmi*, *M. pudendotectus*, *M. confusus* and *M. asymmetricus* (Fig. 2). *M. apri* and *M. salmi* had prevalences of 75-100% and 63-100%, respectively, depending on the age of the animals, whereas *M. asymmetricus* was the rarest with 13-42% (Fig. 3). Differences in species composition between the age groups of the animals were, however, not significant. A significant positive correlation between infections with *M. salmi* and *M. pudendotectus* ($p = 0.000$; Pearson's Rho = 0.493) and an almost significant positive correlation between infections with *M. confusus* and *M. apri* ($p = 0.000$; Rho = 0.564) could be observed.


Metastrongylids are of the most important extraintestinal nematode pathogens of wild boar and organic pigs in terms of animal health, due to the wide distribution of the intermediate earthworm host. A high overall prevalence precluded statistically significant differences between age groups, hinting at a continuous reinfection cycle within the enclosure. So far, correlations between infection intensity and body condition could not be established, however, a tendency of lower body weight in animals with high infections was observed. Other studies could show a correlation between poor body condition and lungworm infection [27].

Table 1. Prevalences and infection intensities for adult *Metastrongylus* spp. and excretion rates for coccidia oocysts and eggs of *Metastrongylus*, *Trichuris* and *Globocephalus* and (only in shoats) *Ascaris* in different age groups. SD: standard deviation

Age group	Parasites	Prevalence (%)	Excretion rates (eggs/oocysts per gram of faeces)				
			Minimum	Maximum	Median	Mean	SD
Shoats (<1 year) N=33 (lungs) N = 31 (faecal samples)	<i>Metastrongylus</i> adults	96.8	0	1020	109	195,5	251.6
	Coccidia oocysts	100	450	88,850	11,650	19,579	20,073.8
	<i>Metastrongylus</i> eggs	90.3	50	400	0	52	103.9
	<i>Trichuris</i> eggs	51.5	50	100	0	11	28.8
	<i>Globocephalus</i> eggs	93.5	50	3000	450	544	649.7
	<i>Ascaris</i> eggs	12.9	50	4950	0	265	955.0
Young pigs (1-2 years) N = 8	<i>Metastrongylus</i> adults	100	1	44	63	22,1	17.2
	Coccidia oocysts	100	4,500	151,150	27,025	50,244	12,667.0
	<i>Metastrongylus</i> eggs	87.5	50	100	0	20	35,0
	<i>Trichuris</i> eggs	62.5	50	50	0	10	21.1
	<i>Globocephalus</i> eggs	100	50	1150	125	305	395.4
Adults (>2 years) N = 8	<i>Metastrongylus</i> adults	87.5	0	307	51,5	87,9	107.0
	Coccidia oocysts	100	1,650	37,550	11,850	14,769	57,487.6
	<i>Metastrongylus</i> eggs	87.5	50	100	0	33	51.6
	<i>Trichuris</i> eggs	37.5	50	50	0	17	25.8
	<i>Globocephalus</i> eggs	100	100	1650	575	608	590.3
TOTAL N= 49 (lungs) N = 47 (faecal samples)	<i>Metastrongylus</i> adults	95.8	0	1020	0	1020	149.6
	Coccidia oocysts	100	450	151,150	14,050	23,980	28,979.5
	<i>Metastrongylus</i> eggs	89.4	0	400	0	49	88.8
	<i>Trichuris</i> eggs	51.1	0	100	0	13	26.5
	<i>Globocephalus</i> eggs	95.7	0	3000	400	550	600.3
	<i>Ascaris</i> eggs	8.0	0	4950	0	174	781.6

This may be due to an influence of age, but irrespective of the risk factors, high worm burdens indicate high infection pressure and in the long run this may lead to serious health problems in the observed herd since in enclosures parasite stages may accumulate (in the environment or in intermediate hosts, depending on the life cycle).

Figure 2. Lung alterations according to age groups.



Coproscopy

Coccidia oocysts could be found in all examined samples (Table 1). Five species of *Eimeria* (*E. deblickei*, *E. perminuta*, *E. polita*, *E. scabra* and *E. porci*) were found in a prevalence of 25-100%, while *Cystoisospora suis* was found in 97.9% of the samples, mostly in shoats (Table 2).

Table 2. Coccidia (*Eimeria* spp., *Cystoisospora suis*) according to species and age group.


	Number of samples	<i>E. deblickei</i>	<i>E. perminuta</i>	<i>E. polita</i>	<i>E. scabra</i>	<i>E. porci</i>	<i>C. suis</i>
Shoats	31	31	30	17	31	8	30
Young pigs	8	8	8	4	8	2	8
Adults	8	8	8	4	6	3	8
TOTAL	47	47	46	25	45	13	46

Excretion intensities varied between 450 and 151,150 oocysts per gram of faeces (Table 1). *E. scabra* was found significantly more often in shoats than in the older pigs ($p = 0.007$) while no correlation could be found for the other coccidia species. At

the same time, *E. scabra* excretion rates were also negatively correlated with body weight ($p = 0.007$; $Rho = -0.446$), which could be an effect of age distribution, although correlations between parasite excretion and age could not be established, probably due to low numbers of animals in some age groups.

In contrast to earlier studies [7, 23] in the present study *E. perminuta* and not *E. suis* was detected.

Figure 3. Prevalences of adult *Metastrongylus* species in different age groups.



Interestingly, all age groups shed considerable amounts of *C. suis* oocysts, which is in contrast to observations in domestic pigs where infection is mostly seen in suckling piglets [24].

Besides eggs of *Metastrongylus*, eggs of *Globocephalus*, *Trichuris* and *Ascaris* could be detected by flotation. Age-related differences were seen only for *Trichuris* eggs which were found mainly in young pigs (62.5%) and *Ascaris* eggs which were

found only in shoats (Table 1). The highest excretion rates were seen for *Ascaris* with 4,950 eggs per gram of faeces (epg) and for *Globocephalus* with 3,000 epg, both in shoats. *Metastrongylus* and *Trichuris* eggs were excreted with a maximum of 400 and 100 epg (Table 1). Quantitative excretion of metastrongylid eggs was positively correlated with the excretion of hookworm eggs ($p = 0.000$; $Rho = 0.679$).

Ondrejková et al. [25] investigated the effect of endoparasites on the oral vaccination against classical swine fever in wild boars and found that parasitic infections influence the efficacy of oral vaccination against swine fever and support the ability of the virus to reproduce and cause disease, and to infect the surrounding the wild boar populations [25]. It must therefore be assumed that parasitic infections render wild boar more prone to other infections.

Conclusion

The rate of lungworm infections of the examined animals was very high in all age groups, as were the infection rates for coccidian. The worm burden and oocyst excretion rates especially in shoats indicated a high infection pressure in young animals which was partially reduced in older boars, probably due to frequent reinfection in combination with induction of immunity [19]. High excretion rates for hookworm, ascarid and whipworm eggs in younger animals also indicate a development of immunity; however, this appears to be slow and incomplete except for *A. suum*, which was excreted only by shoats. The lung alterations, especially emphysema and purulent pneumonia, indicated that animals were more or less severely affected by lungworms (and possibly secondary bacterial infections) in all age groups. Under these circumstances both the high extensity and the high parasite burden indicate that animal health is most likely compromised by parasitic infection since animals constantly remain in the same contaminated environment. Contrary to this, Popiolek et al. observed a higher prevalence of *Metastrongylus* in a wild population than in farmed wild boar [26], so management of farmed wild boar may influence the parasite burden. In the longer term, it is expected that animals with reduced pulmonary function will suffer from reduced food intake, poor growth or even weight loss, as indicated by the approximately 20% of the hunted animals which were in poor body condition, with the consequence of a weakened immune system.

Means of improving animal health in a game enclosure are restricted to controlled hunting to reduce population density, while improved hygiene of feeding areas might be more difficult to implement. Navarro et al. [28] showed that it is important to restrict supplemental feeding because of the high density of wild boar in feeding grounds. They also observed that wild boars defecate before entering feeders which lead to an accumulation of parasite stages, and Nagy et al. [29] reported

an increased density of earthworms infected with metastrongylids in the feeding area of wild boar, so that an accumulation of parasite stages at such sites (where their final hosts aggregate) must be assumed.

The ultimate aim of all measures must be to achieve a balance between game animals and their enclosed habitat to maintain a functional ecological cycle.

Acknowledgements and Conflict of Interest

The authors declare that they have no conflict of interest.

References

1. Barutzki D, Schoirer R, Gothe R. Helminth infections in wild boars kept in enclosures in southern Germany: severity of infections and fecal intensity.[Helmintheninfektionen von Wildschweinen bei Gatterhaltung in Süddeutschland: Befallsstärke und Ausscheidungsintensität]. Tierärztl Prax 1991, 19 (6): 644-648.
2. Acevedoa P, Quirós-Fernández F, Casal J, Vicente J. Spatial distribution of wild boar population abundance: Basic information for spatial epidemiology and wildlife management. Ecol Indic 2014, 36: 594-600.
3. Lindsay D S, Current W L, Taylor J R. Effects of experimentally induced *Isospora suis* infection on morbidity, mortality, and weight gains in nursing pigs. Am J Vet Res 1985, 46 (7):1511-1512.
4. Lindsay D, Neiger R, Hildreth M. Porcine enteritis associated with *Eimeria spinosa* (Henry, 1931) infection. J Parasitol 2002, 88(6):1262-1263.
5. Dauschies A, Imarom S, Ganter M, Bollwahn W. Prevalence of *Eimeria* spp. in sows at piglet producing farms in Germany. J Vet Med B 2004, 51(3): 135-139.
6. Joachim A, Schwarz L, Hinney B, Ruttowski B, Vogl C et al. Which factors influence the outcome of experimental infection with *Cystoisospora suis*? Parasitol Res. 2014, 113(5): 1863-1873.
7. Löwenstein M, Kutzer E. The observation of coccidia in swine. [Studie zur Kenntnis der Schweinekokzidien]. Angew Parasitol. 1989, 30: 117-126.
8. Dold C, Holland C. *Ascaris* and ascariasis. Microbes Infect. 2011, 13(7): 632-637.
9. Andreasen A, Petersen H, Kringel H, Iburg T, Skovgaard K et

- al. Immune and inflammatory response in pigs infected with *Trichuris suis* and *Oesophagostomum dentatum*. *Vet Parasitol*. 2015, 207(3-4): 249-258.
10. Takács A. Contribution to the helminth fauna of wild boar (*Sus scrofa*) in Hungary. [Ein Beitrag zur Helminthenfauna des Wildschweines (*Sus scrofa* L.) in Ungarn]. *Wien. Tierärztl. Mschr*. 1997, 84: 314-316.
11. Wolkers J, Wensing T, Groot Bruinderink G, Schonewille J. Lungworm and stomach worm infection in relation to body fat reserves and blood composition in wild boar (*Sus scrofa*). *Vet Quart* 1994, 16(4): 193-195.
12. Gassó D, Rossi L, Mentaberre G, Casas E, Velarde R et al. An identification key for the five most common species of *Metastrongylus*. *Parasitol Res* 2014, 113(9): 3495-3500.
13. Holló F. Investigation on metastrongylosis in swine: frequency of metastrongylosis and species causing it in Hungary. *Acta Vet Acad Sci Hung*. 1965, 15: 45-60.
14. Alcaide M, Frontera E, Rodríguez M, Saénz I, Domínguez-Alpízar J et al. Pulmonary parasitosis of the iberian pig. metastrongylosis in Spain. [Parasitosis pulmonares del cerdo ibérico: situación actual de la metastrongilosis en España]. *Mundo Ganadero* 2005, 176: 40-44.
15. Holló F. Investigations on metastrongylosis in swine: some remarks to the morphology of *Metastrongylus salmi*, Gedoelst, 1923 with special reference to its differentiation from *M. apri*. *Acta Vet Acad Sci Hung* 1965, 15(3): 259-268.
16. Leignel V, Humbert J, Elard L. Study by ribosomal DNA ITS 2 Sequencing and RAPD analysis on the systematics of four *Metastrongylus* species (Nematoda: Metastrongyloidea). *J Parasitol*. 1997, 83(4): 606-611.
17. Byung-Il Y, Hyeon-Cheol K, Jong-Taek K. Lung worm (*Metastrongylus elongatus*) infection in wild boars (*Sus scrofa*) of the demilitarized zone, Korea. *J Wildl Dis*. 2010, 46(3): 1052-1054.
18. Nosal P, Kowal J, Nowosad B. Structure of Metastrongyloidea in wild boars from southern Poland. *Helminthologia*. 2010, 47(4): 212-218.
19. García-González Á, Pérez-Martín J, Gamito-Santos J, Calero-Bernal R, Alonso M et al. Epidemiologic study of lung parasites (*Metastrongylus* spp.) in wild boar (*Sus scrofa*) in south-western Spain. *J Wildl Dis*. 2013, 49(1):157-162.
20. Joachim A. Helminthosen des Schweins. In: Schnieder T. (Ed.): *Veterinärmedizinische Parasitologie* 2006, Paul Parey, Stuttgart, 6th ed., pp. 369-397.
21. Bauer C. Untersuchungsmethoden. In: Schnieder T. (Ed.): *Veterinärmedizinische Parasitologie* 2006, Paul Parey, Stuttgart, 6th ed.. pp. 84-104.
22. Dausgschies A. Protozoeninfektionen des Schweins. In: Schnieder T. (Ed.): *Veterinärmedizinische Parasitologie* 2006, Paul Parey, Stuttgart, 6th ed., pp. 359-367.
23. Kutzer E. The coccidia of pigs in Austria [Über die Kokzidien des Schweines in Österreich]. *Ztschr Parasitenk* 1960,19: 541-547.
24. Worliczek H, Mundt H, Ruttowski B, Joachim A. Age, not infection dose, determines the outcome of *Isospora suis* infections in suckling piglets. *Parasitol Res* 2009, 105: 157-162.
25. Ondrejková A.: Does the infection with endoparasites influence the effect of oral vaccination against classical swine fever in wild boar? *Acta Vet Brno* 2015, 225-230.
26. Popiolek M. Knecht D, Szczesna-Staskiewicz J, Czerwinska-Rozalow A. Helminths of the wild boar (*Sus scrofa* L.) in natural and breeding conditions. *Bull Vet Inst Pulawy* 2010, 51: 161-166.
27. Syrälä P, Oksanen A, Hälli O, Peltoniemi O, Heinonen M. *Metastrongylus* spp. infection in a farmed wild boar (*Sus scrofa*) in Finland. *Acta Vet Scand* 2010, 51(1): 21.
28. Navarro-Gonzalez N., Fernández-Llario P, Pérez-Martín J., Mentaberre G., López-Martín J et al. Supplemental feeding drives endoparasitic infection in wild boar in Western Spain. *Vet Parasitol* 2013, 196(1-2): 114-123.
29. Nagy G,Csivincsik A, Sugár L. Wild boar density drives *Metastrongylus* infection in earthworm. *Acta Parasitol* 2014, 60(1): 35-39.